


CLAIMS

1. A compound of Formula (I) and pharmaceutically acceptable salts thereof:

5 Formula (I)



10 , wherein: Ar is a nitrogen-containing heteroaromatic ring group selected from a set of groups consisting of a pyridyl group, a pyrimidinyl group, a pyradinyl group, a pyridazinyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyrazolyl group, a pyrrolyl group, an imidazolyl group, an indolyl group, an isoindolyl group, a quinolyl group, an isoquinolyl group, a benzothiazolyl group, and a benzoxazolyl group, which:

15 1) may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group,

20 a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino


25 lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower

10031755-040202

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

alkylcarbamoyloxy group, a di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfanyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or groups represented by a formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p is any of a hydrogen atom, or a lower alkyl group, a lower alkenyl group or a lower alkynyl group which may be substituted with one to three of said substituent(s), or a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolizinyll group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyll group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or


- an aliphatic heterocyclic group selected from a set of groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a tetrahydropyranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, each of which cyclic group may be substituted with one to three of said substituent(s) or, furthermore, may have a bicyclic or tricyclic fused ring of a partial structure selected from a set of groups consisting of:



- ; W_1 is a single bond, an oxygen atom, a sulfur atom, SO , SO_2 , NR_q , SO_2NR_q , $N(R_q)SO_2NR_r$, $N(R_q)SO_2$, $CH(OR_q)$, $CONR_q$, $N(R_q)CO$, $N(R_q)CONR_r$, $N(R_q)COO$, $N(R_q)CSO$, $N(R_q)COS$, $C(R_q)=CR_r$, $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_q$, $OC(S)NR_q$, $SC(O)$, $SC(O)NR_q$ and $C(O)O$ (wherein: R_q and R_r are each either a substituent selected from a set of groups consisting of (i) a hydrogen atom, (ii) a substituent selected from a set of groups consisting of a lower alkyl group, a cyclo lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, a lower alkoxy group, a lower

10031755.040202

- alkoxycarbonyl group, a lower alkoxy-carbonylamino group, a lower alkoxy-carbonylamino lower alkyl group, a lower alkyl-carbamoyl group, a di-lower alkyl-carbamoyl group, a carbamoyloxy group, a lower alkyl-carbamoyloxy group, di-
- 5 lower alkyl-carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a
- 10 lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or (iii) a lower alkyl group, an aryl
- 15 group or an aralkyl group which may be substituted with one to three of said substituent(s).); Y_1 and Y_2 are each, the same or different, a single bond or a straight-chain or branched lower alkylene group which may have any of said bicyclic or tricyclic fused ring);
- 20 2) may have a five- to seven-membered fused ring selected from a set of groups consisting of:




- which may be formed together with the carbon atom of said nitrogen-containing heteroaromatic cyclic group, on which
- 25 the substituent, which is selected from a set of groups

- consisting of a lower alkyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group,
- 5 lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group,
- 10 an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group,
- 15 an aroylamino group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, and a lower alkanoylamidino lower alkyl group (hereinafter indicated as ring-substituent) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom
- 20 and/or a nitrogen atom on said ring-substituent;

or,

3) may have a five- to seven-membered ring selected from a set of groups consisting of:



- 25 which may be formed together with the carbon atom of said

nitrogen-containing heteroaromatic group on which a substituent represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein: Y_1 , W_1 , Y_2 and R_p have the same meanings as stated above) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent.

; X and Z are each, the same or different, a carbon atom or a nitrogen atom, or being taken together with R_1 or R_2 and/or R_3 which may exist on X or Z, form a CH or a nitrogen atom; Y is CO, SO or SO_2 ; R_1 is any of a hydrogen atom or a substituent represented by a formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s is any of a hydrogen atom or a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a cyclo lower alkyl group, an aryl group, and a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolizinyll group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyll group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- an aliphatic heterocyclic group selected from a set of groups consisting of an isoxazoliny1 group, an isoxazolidiny1 group, a tetrahydropyridy1 group, an imidazolidiny1 group, a tetrahydrofurany1 group, a
- 5 piperaziny1 group, a piperidiny1 group, a pyrrolidiny1 group, pyrroliny1 group, a morpholino group, a tetrahydroquinoliny1 group and a tetrahydroisoquinoliny1 group, all of which may be substituted with one to three of said substituent(s); W_2 is a single bond, an oxygen atom, a
- 10 sulfur atom, SO , SO_2 , NR_t , SO_2NR_u , $N(R_t)SO_2NR_u$, $N(R_t)SO_2$, $CH(OR_t)$, $CONR_t$, $N(R_t)CO$, $N(R_t)CONR_u$, $N(R_t)COO$, $N(R_t)CSO$, $N(R_t)COS$, $C(R_v)=CR_r$, $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_t$, $OC(S)NR_t$, $SC(O)$, $SC(O)NR_t$ and $C(O)O$ (wherein: R_t and R_u are each a hydrogen atom or a substituent selected from a set of
- 15 groups consisting of a lower alkyl group, a hydroxy group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group,
- 20 a carboxy lower alkyl group, a carbamoyl lower alkyl group, a lower alkoxy group, a lower alkoxy carbonyl group, a lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower
- 25 alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower


11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of said substituent(s)); Y_3 and Y_4 are each, the same or different, a single bond or a straight-chain or branched lower alkylene group), or R_1 is an lower alkyl group which may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl

group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a substituent selected from groups represented by the formula $Y_3-W_2-Y_4-R_8$ (wherein: R_8 , W_2 , Y_3 and Y_4 have the same meanings as stated above), or R_1 forms a nitrogen atom together with X.); R_2 and R_3 are each independently, the same or different, a hydrogen atom, a hydroxy group, a lower alkyl group, a lower alkoxy group, or a substituent represented by the formula $Y_3-W_2-Y_4-R_8$ (wherein: R_8 , W_2 , Y_3 and Y_4 have the same meanings as stated above), or one of R_2 or R_3 forms, together with R_1 and X, a saturated five- to eight-membered cyclic group selected from sets of groups consisting of (a) and (b):




15 and




and another one of R_2 or R_3 binds to a carbon atom or a nitrogen atom on the ring, or to a carbon atom, an oxygen atom and/or nitrogen atom on said ring-substituent of said ring to form a five- to seven-membered ring, or R_2 and R_3 are combined to form a spiro cyclo lower alkyl group, or R_2 and R_3 are combined, together with Z on which they exist to form an oxo (keto, or carbonyl) group, or R_2 and R_3 form, together with Z, R_1 and X, on which they stand, a saturated or an unsaturated five- to eight membered cyclic group

which may be selected from sets of groups of (a) and (b):



and



- 5 , which may contain one or more kinds of hetero atom(s) selected from a group of a nitrogen atom, an oxygen atom and a sulfur atom, and which may be fused with any of a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an
- 10 imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolydinyl group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a
- 15 pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a
- 20 benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group(s) selected from a set of groups consisting of an isoxazoliny group, an isoxazolidiny group, a tetrahydropyridyl group, an imidazolidiny group, a tetrahydrofuranyl group, a tetrahydropyranyl group, a piperaziny group, a piperidinyl group, a pyrrolidinyl group, pyrroliny group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, which may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a spiro cyclo lower alkyl group which may be substituted, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower


1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25

alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and a substituent selected from groups
 5 represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p , W_1 , Y_1 and Y_2 have the same meanings as stated above); R_4 and R_5 are each, the same or different, a hydrogen atom, halogen atoms, a hydroxy group, an amino group, or a substituent represented by the formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s , W_2 , Y_3
 10 and Y_4 have the same meanings as stated above), or any of a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of the same or different substituent(s) selected from both a set of groups consisting of a lower alkyl group, a cyano group, a nitro
 15 group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower
 20 alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower
 25 alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower

- alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and groups represented by the formula
- 5 $Y_3-W_2-Y_4-R_5$ (wherein: R_5 , W_2 , Y_3 and Y_4 have the same meanings as stated above); and the formula \equiv represents either a single bond or a double bond.


2. A compound according to claim 1, having a structure of
- 10 Formula (I-a), and pharmaceutically acceptable salts thereof:

Formula (I-a)




- , wherein: Ar_a is a nitrogen-containing heteroaromatic ring
- 15 group selected from a set of groups consisting of a pyridyl group, a pyrimidinyl group, a pyradinyl group, a pyridazinyl group, a thiazolyl group, a pyrazolyl group, and an imidazolyl group, and said nitrogen-containing heteroaromatic ring group
- 20 1') may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, halogen atoms, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a
- 25 halo lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower

FULL-SCALE



- heteroaromatic ring group, on which a substituent selected from a set of groups consisting of a lower alkyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a halo lower alkyl group, a carbamoyl lower alkyl group, a lower alkoxy group, a lower alkoxy carbonyl group, a lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a lower alkyl carbamoyloxy group, a lower alkylamino group, di-lower alkylamino group, an amino lower alkyl group, a lower alkylamino lower alkyl group, di-lower alkylamino lower alkyl group, a lower alkanoylamino group, and an aroylamino group exists, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom, each of which exists in said ring-substituent(s);
- or,
- 3') may form a five- to six-membered ring selected from a set of groups consisting of:




- together with a carbon atom on said nitrogen-containing heteroaromatic ring group, on which a substituent represented by the formula $Y_{1a}-W_{1a}-Y_{2a}-R_{pa}$ (wherein: Y_{1a} , W_{1a} , Y_{2a} and R_{pa} have the same meanings as stated above) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom in said ring-substituent(s); X_a and Z_a are each, the same or different, a carbon atom or a nitrogen atom, or optionally being taken together with R_{1a} or R_{2a} and/or R_{3a} on them form a CH or a nitrogen atom; Y_a is a CO, SO or SO_2 , R_{1a} is a hydrogen atom

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- or a substituent represented by a formula $Y_{3a}-W_{2a}-Y_{4a}-R_{sa}$ (wherein: R_{sa} is a hydrogen atom or a lower alkyl group, a lower alkenyl group, a cyclo lower alkyl group, an aryl group, or a heteroaromatic ring group selected from a group
- 5 consisting of an indolyl group, or an aliphatic heterocyclic group selected from a group of a tetrahydropyridyl group, a piperadiny group, a piperidinyl group, a pyrrolidinyl group and a morpholino group, all of which groups may be substituted with one to three of the
- 10 same or different said substituent(s); W_{2a} is a single bond, NR_{ta} , $CH(OR_{ta})$, $CONR_{ta}$, $N(R_{ta})CO$, $N(R_{ta})COO$, $OC(O)NR_{ta}$ or $C(O)O$ (wherein: R_{ta} is a hydrogen atom, a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of said substituent(s)); Y_{3a}
- 15 and Y_{4a} are each, the same or different, a single bond, or a straight-chain or branched lower alkylene group); or R_1 is a lower alkyl group which may be substituted with one to three substituent(s) selected from both a set of groups consisting of a lower alkyl group, a hydroxyl group, a
- 20 carbamoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a lower alkoxy group, a lower alkoxy carbonyl group, a lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a
- 25 carbamoyloxy group, a lower alkylcarbamoyloxy group, a lower alkylamino group, a di-lower alkylamino group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a lower alkanoylamino group, and an aroylamino group, and groups


- represented by the formula $Y_{3a}-W_{2a}-Y_{4a}-R_{sa}$ (wherein: R_{sa} , W_{2a} , Y_{3a} and Y_{4a} have the same meanings as stated above), or form a nitrogen atom, together with X; R_{2a} and R_{3a} are each independently, the same or different, a hydrogen atom, or a substituent of a hydroxy group, a lower alkyl group, a lower alkoxy group, or the one represented by the formula $Y_{3a}-W_{2a}-Y_{4a}-R_{sa}$ (wherein: R_{sa} , W_{2a} , Y_{3a} and Y_{4a} have the same meanings as stated above), or any one of R_{2a} or R_{3a} forms, together with R_{1a} and X_a , a saturated five- to eight-membered cyclic group selected from a set of groups consisting of (a-1) and (b-1),

(a-1)





and


(b-1)



- and the other one binds to a carbon atom or a nitrogen atom on the ring, or to a carbon atom, an oxygen atom and/or nitrogen atom on said ring-substituent to form a five- to seven-membered ring, or R_{2a} and R_{3a} are combined to form a spiro cyclo lower alkyl group, or R_{2a} and R_{3a} are combined with Z on which they stand to form an oxo (a keto, or carbonyl) group, or R_{2a} and R_{3a} form, together with Z_a on which they stand, R_{1a} and X_a , a saturated or an unsaturated five- to eight membered cyclic group which may be selected from sets of groups of (a-1) and (a-2):



and




- which may have one or more kinds of hetero atom(s), and
- 5 which may be substituted with one to three of the same or different substituent(s) selected both from a set of groups consisting of a lower alkyl group, a spiro cyclo lower alkyl group which may be substituted, a hydroxy group, a hydroxy lower alkyl group, lower alkoxy group, a lower
- 10 alkoxycarbonyl group, a lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a lower alkylcarbamoyloxy group, a lower alkylamino group, a di-lower alkylamino group, an amino lower alkyl group, a lower alkylamino lower alkyl
- 15 group, a di-lower alkylamino lower alkyl group, a lower alkanoylamino group and an aroylamino group, and groups represented by the formula $Y_{1a}-W_{1a}-Y_{2a}-R_{pa}$ (wherein: R_{pa} , W_{1a} , Y_{1a} and Y_{2a} have the same meanings as stated above), and, furthermore, which may be fused with a cyclo lower alkyl
- 20 group, an aryl group, a heteroaromatic ring group selected from a group of a pyridyl group and a pyrazolyl group, and an aliphatic heterocyclic group selected from a group of piperidinyl group and a pyrrolidinyl group; R_{4a} and R_{5a} are each, the same or different, a hydrogen atom or a

substituent consisting of halogen atoms, a hydroxy group, an amino group, or the one represented by the formula $Y_{3a}-W_{2a}-Y_{4a}-R_{sa}$ (wherein: R_{sa} , W_{2a} , Y_{3a} and Y_{4a} have the same meanings as stated above), or a lower alkyl group, an aryl group or an aralkyl group, each of which may be substituted with one to three of the same or different substituent(s) selected from both a set of groups consisting of a lower alkyl group, a hydroxy lower alkyl group, a halo lower alkyl group, a lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a lower alkylamino group, a lower alkylamino lower alkyl group, a lower alkanoylamino group, and an aroylamino group, and groups represented by the formula $Y_{3a}-W_{2a}-Y_{4a}-R_{sa}$ (wherein: R_{sa} , W_{2a} , Y_{3a} and Y_{4a} have the same meanings as stated above); and the formula \sim is a single bond or a double bond.

3. A compound according to claim 1 and 2, having a structure of Formula (I-b) and pharmaceutically acceptable salts thereof,


Formula (I-b)



wherein: Ar_b is a nitrogen-containing heteroaromatic ring group selected from a set of groups comprising a pyridyl group and a pyrazolyl group, which:


1'') may be substituted with one to three substituent(s)

selected from both a set of groups consisting of a hydroxy group, halogen atoms, a lower alkanoyloxy group, a hydroxy lower alkyl group, a lower alkoxy group, a lower alkoxycarbonyl group, an amino group, and a lower alkylamino lower alkyl group, and groups represented by a formula $Y_{1b}-W_{1b}-Y_{2b}-R_{pb}$ (wherein: R_{pb} is a hydrogen atom or a lower alkyl group, a lower alkenyl group or a lower alkynyl group which may be substituted with one to three of said substituent(s), or a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of a pyridyl group and a pyrazolyl group, or an aliphatic heterocyclic group selected from a set of groups consisting of isoxazolinyl group, a tetrahydropyridyl group, a piperadinyl group, a piperidinyl group, a pyrrolidinyl group, a morpholino group and a tetrahydroisoquinolinyl group, each of which cyclic substituent groups may be substituted with one to three of said substituent(s) and, furthermore, may have a bicyclic or tricyclic fused ring, which contains the partial structure of which is selected from a group consisting of:




; W_{1b} is NR_{qb} , $N(R_{qb})SO_2$, $CONR_{qb}$, $N(R_{qb})CO$, $N(R_{qb})COO$, $OC(O)$, and $C(O)O$ (wherein: R_{qb} and R_{rb} is a hydrogen atom, or a substituent selected from a set of groups consisting of a hydroxy group, halogen atoms, a cyclo lower alkyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a lower alkoxy group, a lower alkoxycarbonyl group, an amino group, and a lower alkylamino lower alkyl group, or a lower

- alkyl group, an aryl group or an aralkyl group, which may be substituted with one to three of said substituent(s)); Y_{1b} and Y_{2b} are each, the same or different, a single bond or a straight-chain or branched lower alkylene group which
- 5 may have a said bicyclic or tricyclic fused ring);
- 2'') may have a five- or six-membered ring selected from a group consisting of:




- which is together with the ring carbon atom on which a
- 10 substituent selected from a group consisting of a lower alkanoyloxy group, a hydroxy lower alkyl group, a lower alkoxy group, a lower alkoxy carbonyl group and a lower alkylamino lower alkyl group stands, a carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or
- 15 a nitrogen atom in said ring-substituent; or,
- 3'') may form a five- or six-membered ring selected from a group consisting of:




- which is together with the ring-carbon atom on which a
- 20 substituent represented by the formula $Y_{1b}-W_{1b}-Y_{2b}-R_{pb}$ (wherein: Y_{1b} , W_{1b} , Y_{2b} and R_{pb} have the same meanings as stated above) stands, a carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom in said ring-substituent; X_b and Z_b are each, the same
- 25 or different, a carbon atom or a nitrogen atom, or X_b and Z_b form a CH or a nitrogen atom, being taken together with R_{1b} or R_{2b} and/or R_{3b} on them; Y_b is a CO, SO or SO_2 ; R_{1b} is a

- hydrogen atom or a substituent represented by a formula $Y_{3b}-W_{2b}-Y_{4b}-R_{sb}$ (wherein: R_{sb} is a hydrogen atom or a lower alkyl group, a cyclo lower alkyl group, and an aryl group, which may be substituted with one to three of said
- 5 substituent(s); W_{2b} is a single bond, $N(R_{tb})COO$ or $C(O)O$ (wherein: R_{tb} is a hydrogen atom or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of said substituent(s)); Y_{3b} and Y_{4b} are each, the same or different, a single bond, or a straight-
- 10 chain or branched lower alkylene group), or a lower alkyl group which may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a hydroxy lower alkyl group and the one represented by the formula $Y_{3b}-W_{2b}-Y_{4b}-R_{sb}$ (wherein: R_{sb} ,
- 15 W_{2b} , Y_{3b} and Y_{4b} have the same meanings as stated above), or forms a nitrogen atom, together with X; R_{2b} and R_{3b} are each independently, the same or different, a hydrogen atom, a hydroxy group, a lower alkyl group, a lower alkoxy group, or a substituent represented by the formula $Y_{3b}-W_{2b}-Y_{4b}-R_{sb}$
- 20 (wherein: R_{sb} , W_{2b} , Y_{3b} and Y_{4b} have the same meanings as stated above), or either R_{2b} or R_{3b} forms, together with R_{1b} and X_b , a saturated five- to eight-membered cyclic group selected from sets of groups of (b-1) and (b-2),




25 and



and the other one binds to a carbon atom or a nitrogen atom


on the ring, or to a carbon atom, an oxygen atom and/or nitrogen atom on said ring-substituent to form a five- to seven-membered ring, or R_{2b} and R_{3b} are combined to form a spiro cyclo lower alkyl group, or they (R_{2b} and R_{3b}) are combined furthermore with Z on which they stand to form an oxo (a keto, or carbonyl) group, or they (R_{2b} and R_{3b}) form, together with Z_b on which they stand, R_{1b} and X_b , a saturated or an unsaturated five- to seven-membered cyclic group which may be selected from sets of groups of (b-1) and (b-2):

(b-1)



and

(b-2)




which may either have one or more kinds of hetero atom(s) selected from a group of a nitrogen atom, an oxygen atom and a sulfur atom, or which may be fused with a cyclo lower alkyl group, an aryl group and an aliphatic heterocyclic group selected from a group of a piperidinyl group and a pyrrolidinyl group, all of which cyclic groups may be substituted with one to three of the same or different substituent(s) selected both from a set of groups consisting of a lower alkyl group, a spiro cyclo lower alkyl group which may be substituted, a hydroxy lower alkyl group and a lower alkoxy carbonyl group, and groups represented by the formula $Y_{1b}-W_{1b}-Y_{2b}-R_{pb}$ (wherein: R_{pb} , W_{1b} , Y_{1b} and Y_{2b} have the same meanings as stated above); R_{4b} and R_{5b} are each independently, the same or different,

- or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of substituents comprising any of a hydrogen atom, halogen atoms or a substituent represented by the formula $Y_{3b}-W_{2b}-Y_{4b}-R_{5b}$ (wherein: R_{5b} , W_{2b} , Y_{3b} and Y_{4b} have the same meanings as stated above), or a substituent selected from a set of groups consisting of a lower alkyl group, a hydroxy lower alkyl group, a halo lower alkyl group, a lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a lower alkylamino group, a lower alkylamino lower alkyl group, a lower alkanoylamino group, and an aroylamino group.; and the formula $---$ means a single bond or a double bond.

4. A compound according to any one of claim 1 to claim 3, having a structure of Formula (I-p) and pharmaceutically acceptable salts thereof,

Formula (I-p)



- wherein: Ar_p is a nitrogen-containing heteroaromatic ring group which may be substituted, X_p is a carbon atom (CH) or a nitrogen atom, R_{1p} is a hydrogen atom or a lower alkyl group which may be substituted, R_{2p} is a hydrogen or an oxo group (which forms carbonyl group, together with the carbon

atom on which it stands), or forms, together with the carbon atom on which it stands, R_{1p} and X_p , a saturated or an unsaturated five- or six-membered cyclic group which may have one or more kinds of hetero atom(s) selected from a group of a nitrogen atom and a sulfur atom or which may be substituted; R_{4p} and R_{5p} are each, the same or different, any of a hydrogen atom, halogen atoms, a hydroxy group, an amino group or a lower alkyl group, an aryl group or an aralkyl group which may be substituted.

10

5. A compound according to claim 1, wherein the compound is $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(2-\text{octylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(2-\text{methyl-4,4-dimethylpentylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(5-\text{methoxyindan-2-ylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(2-\text{methylindan-2-ylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(5-\text{chloroindan-2-ylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(6-\text{methylpyridin-2-yl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(\text{pyrrolidin-2-yl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(\text{t-butylaminomethyl})\text{pyrazol-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(\text{pyrazolo}[5,4-b]\text{pyridin-3-yl})\text{urea}$, $N'-(\text{pyrrolidino}[2,1-b]\text{isoindolin-4-on-8-yl})-N-(5-(1-\text{hydroxymethylcyclopentylaminomethyl})\text{pyrazol-3-yl})\text{urea}$,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- N'-(pyrrolidino[2,1-b]-4-oxoisindolin-8-yl)-N-(5-(N-t-butyl-N-methyl-aminomethyl)pyrazol-3-yl)urea, N'-
(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-1,2,5,6-tetrahydropyridin-4-yl)pyridin-2-yl)urea, N'-
5 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-4-piperidyl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)urea, N'-
(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-3-piperidyl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-b]-4-oxoisindolin-8-yl)-N-(4-(1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-acetyl-3-piperidyl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(piperidino[3,4-c]pyridin-5-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(pyrrolidino[3,4-c]pyridin-5-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(cyclohexylaminoethyl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-cyclohexylpyrrolidin-3-yl)pyridin-2-yl)urea (compound 180), N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzylpyrrolidin-3-yl)pyridin-2-yl)urea, N'-(N-cyclopentyl-3-methylisoindolin-1-on-4-yl)-N-(pyridin-2-yl)urea, N'-(3-t-butylisoindolino[3,2-b]oxazolidin-4-on-8-yl)-N-(4-(N-benzylpyrrolidin-3-yl)pyridin-2-yl)urea, N'-(2-methylisoindolino[3,2-b]perhydro-1,3-oxazin-5-on-9-yl)-N-(4-(N-benzylpyrrolidin-3-yl)pyridin-2-yl)urea, or N'-(isoindolino[2,3-b]perhydro-1,4-methano-6,11a-benzoxazin-11-on-7-yl)-N-(pyridin-2-yl)urea.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

6. A compound according to claim 1, wherein the compound is
 N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(5-(2-
 octylaminomethyl)pyrazol-3-yl)urea, N'-(pyrrolidino[2,1-
 5 b]isoindolin-4-on-8-yl)-N-(5-(2-methyl-4,4-
 dimethylpentylaminomethyl)pyrazol-3-yl)urea, N'-
 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(5-(5-
 methoxyindan-2-ylaminomethyl)pyrazol-3-yl)urea,
 N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(5-(2-
 10 methylindan-2-ylaminomethyl)pyrazol-3-yl)urea, N'-
 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(5-(5-
 chloroindan-2-ylaminomethyl)pyrazol-3-yl)urea, N'-
 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-
 1,2,5,6-tetrahydropyridin-4-yl)pyridin-2-yl)urea, N'-
 15 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-benzyl-4-
 piperidyl)pyridin-2-yl)urea, N'-(pyrrolidino[2,1-
 b]isoindolin-4-on-8-yl)-N-(piperidino[3,4-c]pyridin-6-
 yl)urea, N'-(pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-
 (N-cyclohexylpyrrolidin-3-yl)pyridin-2-yl)urea, N'-
 20 (pyrrolidino[2,1-b]isoindolin-4-on-8-yl)-N-(4-(N-
 benzylpyrrolidin-3-yl)pyridin-2-yl)urea, N'-(3-t-
 butylisoindolino[3,2-b]oxazolidin-4-on-8-yl)-N-(4-(N-

benzylpyrrolidin-3-yl)pyridin-2-yl)urea, N'-(2-methylisoidolino[3,2-b]perhydro-1,3-oxazin-5-on-9-yl)-N-(4-(N-benzylpyrrolidin-3-yl)pyridin-2-yl)urea, or N'-(isoidolino[2,3-b]perhydro-1,4-methano-6,11a-benzoxazin-11-on-7-yl)-N-(pyridin-2-yl)urea.

7. A method of manufacturing a compounds of Formula (I) and pharmaceutically acceptable salts thereof characterized by reacting compounds of Formula (III) with a compound of Formula (IV):

Formula (III)



wherein: X and Z are each, the same or different, a carbon atom or a nitrogen atom, or a CH or a nitrogen atom, together with R₁₀ or R₂₀ and/or R₃₀ which bind to X or Z; Y is a CO, SO or SO₂; R₁₀ is a hydrogen atom or a substituent represented by a formula Y₃₀-W₂₀-Y₄₀-R₅₀ (wherein: R₅₀ is a hydrogen atom or a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolydiny group, an

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisindolyl
- 5 group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl
- 10 group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group(s) selected from a set of groups consisting of an isoxazoliny group, an isoxazolidinyl group, a tetrahydropyridyl group, an
- 15 imidazolidinyl group, a tetrahydrofuranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group which may be substituted with one to three of said
- 20 substituents; W_{20} is a single bond, an oxygen atom, a sulfur atom, SO , SO_2 , NR_{t0} , SO_2NR_{t0} , $N(R_{t0})SO_2NR_{u0}$, $N(R_{t0})SO_2$, $CH(OR_{t0})$, $CONR_{t0}$, $N(R_{t0})CO$, $N(R_{t0})CONR_{u0}$, $N(R_{t0})COO$, $N(R_{t0})CSO$, $N(R_{t0})COS$, $C(R_{v0})=CR_{t0}$, $C \equiv C$, CO , CS , $OC(O)$, $OC(O)NR_{t0}$, $OC(S)NR_{t0}$, $SC(O)$, $SC(O)NR_{t0}$ and $C(O)O$ (wherein: R_{t0} and R_{u0}
- 25 are each a hydrogen atom or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group which may be protected, a cyano group, halogen atoms, a nitro


1-
2-
3-
4-
5-
6-
7-
8-
9-
10-
11-
12-
13-
14-
15-
16-
17-
18-
19-
20-
21-
22-
23-
24-
25-
26-
27-
28-
29-
30-
31-
32-
33-
34-
35-
36-
37-
38-
39-
40-
41-
42-
43-
44-
45-
46-
47-
48-
49-
50-
51-
52-
53-
54-
55-
56-
57-
58-
59-
60-
61-
62-
63-
64-
65-
66-
67-
68-
69-
70-
71-
72-
73-
74-
75-
76-
77-
78-
79-
80-
81-
82-
83-
84-
85-
86-
87-
88-
89-
90-
91-
92-
93-
94-
95-
96-
97-
98-
99-
100-

- group, a carboxyl group which may be protected, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group which may be protected, a cyano lower alkyl group, a halo lower alkyl
- 5 group, a carboxy lower alkyl group which may be protected, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a
- 10 carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group which may be protected, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group which may be protected, a lower alkylamino lower
- 15 alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower
- 20 alkylsulfonylamino group, a hydroxyimino group which may be protected, and a lower alkoxyimino group, or said substituent(s)); Y_{30} and Y_{40} are each, the same or different, a single bond or a straight-chain or branched lower alkylene group); or a lower alkyl group which may be
- 25 substituted with one to three of substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group which may be protected, a cyano group, halogen atoms, a nitro group, a carboxyl group which may be protected, a carbamoyl group, a formyl group, a lower


10034756-040222

- alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group which may be protected, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group which may be protected, a carbamoyl lower alkyl group,
- 5 lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group,
- 10 an amino group which may be protected, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a
- 15 lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group which may be protected, and a lower alkoxyimino group, or a (set of
- 20 substituents) substituent represented by the formula $Y_{30}-W_{20}-Y_{40}-R_{50}$ (wherein: R_{50} , W_{20} , Y_{30} and Y_{40} have the same meanings as stated above), or R_{10} forms a nitrogen atom, together with X; R_{20} and R_{30} are each independently, the same or different, any of a hydrogen atom, a hydroxy group
- 25 which may be protected, a lower alkyl group, a lower alkoxy group, or a substituent represented by the formula $Y_{30}-W_{20}-Y_{40}-R_{50}$ (wherein: R_{50} , W_{20} , Y_{30} and Y_{40} have the same meanings as stated above), or either one of R_{20} and R_{30} forms, together with R_{10} and X, a saturated five- to eight-


membered cyclic group selected from a sets of groups consisting of (a) and (b):



5 and




and the other one binds either to a carbon atom or a nitrogen atom on the ring, or to a carbon atom, an oxygen atom and/or a nitrogen atom on the ring-substituent(s) on said ring, to form a five- to seven-membered ring, or R₂₀ and R₃₀ are combined to form a spiro cyclo alkyl group, or to form, together with Z, on which they stand, an oxo (keto, carbonyl) group, or, to form, together with Z on which they stand, R₁₀ and X, a heteroaromatic ring consisting of a saturated or an unsaturated five- to eight-membered cyclic ring selected from sets of groups consisting of (a) and (b)



and


(b)



- ,which may either contain one or more kinds of hetero atoms selected from the group consisting of a nitrogen atom, an oxygen atom and a sulfur atom, or which may be fused with a ring selected from a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolydinyll group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group(s) selected from a set of groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a

protected, and a lower alkoxyimino group, or a set of substituent(s) represented by a formula $Y_{10}-W_{10}-Y_{20}-R_{p0}$ (wherein: R_{p0} is a hydrogen atom, or a lower alkyl group, a lower alkenyl group, or a lower alkynyl group, which may be substituted with one to three of said substituent(s), or a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indoldindinyl group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group(s) selected from a set of groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a tetrahydropyranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, which may be substituted

with one to three of said substituent(s), or, furthermore, may have on it a bicyclic or tricyclic fused ring which contains a partial structure selected from a set of groups comprising:



; W_{10} is a single bond, an oxygen atom, a sulfur atom, SO , SO_2 , NR_{q0} , SO_2NR_{q0} , $N(R_{q0})SO_2NR_{r0}$, $N(R_{q0})SO_2$, $CH(OR_{q0})$, $CONR_{q0}$, $N(R_{q0})CO$, $N(R_{q0})CONR_{q0}$, $N(R_{q0})COO$, $N(R_{q0})CSO$, $N(R_{q0})COS$, $C(R_{q0})=CR_{r0}$, $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_{q0}$, $OC(S)NR_{q0}$, $SC(O)$, $SC(O)NR_{q0}$ and $C(O)O$ (wherein: R_{q0} and R_{r0} are each either a hydrogen atom or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of substituent(s) selected from a set of groups consisting of a lower alkyl group, a cyclo lower alkyl group, a hydroxyl group which may be protected, a cyano group, halogen atoms, a nitro group, a carboxyl group which may be protected, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group which may be protected, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group which may be protected, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group which may be protected, a lower alkylamino group, a di-lower alkylamino group, a tri-lower


100-1751-210-01

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

alkylammonio group, an amino lower alkyl group which may be
 protected, a lower alkylamino lower alkyl group, a di-lower
 alkylamino lower alkyl group, a tri-lower alkylammonio
 lower alkyl group, a lower alkanoylamino group, an
 5 aroylamino group, a lower alkanoylamidino lower alkyl group,
 a lower alkylsulfinyl group, a lower alkylsulfonyl group, a
 lower alkylsulfonylamino group, a hydroxyimino group which
 may be protected, and a lower alkoxyimino group, or from
 said substituent(s)); Y_{10} and Y_{20} are each, the same or
 10 different, a single bond or a straight-chain or branched
 lower alkylene group which may have a bicyclic or tricyclic
 fused ring); R_{40} and R_{50} are each, the same or different,
 either a hydrogen atom, halogen atoms, a hydroxyl which may
 be protected, an amino group which may be protected, or a
 15 lower alkyl group, an aryl group or an aralkyl group which
 may be substituted with one to three of the same or
 different substituents selected from a set of the groups
 consisting of: the one represented by the formula $Y_{30}-W_{20}-$
 $Y_{40}-R_{60}$ (wherein: R_{60} , W_{20} , Y_{30} and Y_{40} have the same meanings
 20 as stated above), the one which may be selected from a set
 of groups consisting of a lower alkyl group, a cyano group,
 a nitro group, a carboxyl group which may be protected, a
 carbamoyl group, a formyl group, a lower alkanoyl group, a
 lower alkanoyloxy group, a hydroxy lower alkyl group which
 25 may be protected, a cyano lower alkyl group, a halo lower
 alkyl group, a carboxy lower alkyl group which may be
 protected, a carbamoyl lower alkyl group, a lower alkoxy
 group, a lower alkoxy carbonyl group, lower
 alkoxy carbonylamino group, a lower alkoxy carbonylamino

lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group which may be protected, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group which may be protected, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group which may be protected, and a lower alkoxyimino group, and the one represented by the formula $Y_{30}-W_{20}-Y_{40}-R_{50}$ (wherein: R_{50} , W_{20} , Y_{30} and Y_{40} have the same meanings as stated above); the Formula --- is a single bond or a double bond, or is made to react with a compound represented by Formula (IV)


Formula (IV)



wherein: Ar_0 is a nitrogen-containing heteroaromatic ring group selected from a set of groups consisting of a pyridyl group, a pyrimidinyl group, a pyradinyl group, a pyridazinyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyrazolyl group, a pyrrolyl group, an imidazolyl group, an indolyl group, an isoindolyl group, a quinolyl group, an isoquinolyl group, a benzothiazolyl group, a benzoxazolyl group, which:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- 1) may be substituted with one to three of the same or different substituent(s) selected from a set of groups a lower alkyl group, a hydroxyl group which may be protected, a cyano group, halogen atoms, a nitro group, a carboxyl group which may be protected, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group which may be protected, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group which may be protected, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group which may be protected, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkyl ammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkyl ammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group which may be protected and a lower alkoxyimino group, or a substituent selected from groups represented by a formula $Y_{10}-W_{10}-Y_{20}-R_{p0}$ (wherein: R_{p0} , W_{10} , Y_{10} and Y_{20} have the same meanings as stated above); or
- 2) may have a five- to seven-membered ring selected from a set of groups consisting of




10031735.013333

- , which may be protected, and together with the carbon atom on the ring on which the substituent selected from a set of groups consisting of a lower alkyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group which may be protected, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group which may be protected, a carbamoyl lower alkyl group, a lower alkoxy group, a lower alkoxy carbonyl group, a lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, a di-lower alkyl carbamoyloxy group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group which may be protected, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkylfulfynyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group and a lower alkanoylamidino lower alkyl group (hereinafter indicated as ring-substituent(s) which may be protected) stands, a carbon atom next to said carbon atom and a carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent(s) which may be protected, all taken together;


or

3) may have a five- to seven-membered ring selected from a set of groups consisting of




5 ,which may be protected, and together with the carbon atom
on the ring on which a substituent selected from groups
represented by the formula of $Y_{10}-W_{10}-Y_{20}-R_{p0}$ (wherein:
 Y_{10}, W_{10}, Y_{20} and R_{p0} have the same meanings as stated above)
stands, a carbon atom next to said carbon atom and a carbon
10 atom, an oxygen atom and/or a nitrogen atom on said ring-
substituent(s) which may be protected, all taken together,
to give a compound of Formula (II)

Formula (II)



15 , wherein Ar₁, X, Y, Z, R₁₀, R₂₀, R₃₀, R₄₀ and the Formula ~~---~~
have the same meanings as stated above, and then, if
necessary, removing the protecting group(s), to give a
compound of Formula (I) according to claim 1 and
pharmaceutically acceptable salts thereof:




- , wherein: Ar is a nitrogen-containing heteroaromatic ring group selected from the groups consisting of a pyridyl group, a pyrimidinyl group, a pyradinyl group, a pyridazinyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyrazolyl group, a pyrrolyl group, an imidazolyl group, an indolyl group, an isoindolyl group, a quinolyl group, an isoquinolyl group, a benzothiazolyl group, and a benzoxazolyl group, and said
- 10 nitrogen-containing heteroaromatic ring group, which:
- 1) may be optionally substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl
- 15 group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower
- 20 alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower
- 25 alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25


di-lower alkylamino lower alkyl group, a tri-lower
alkylammonio lower alkyl group, a lower alkanoylamino group,
an aroylamino group, a lower alkanoylamidino lower alkyl
group, a lower alkylsulfinyl group, a lower alkylsulfonyl
5 group, a lower alkylsulfonylamino group, a hydroxyimino
group and a lower alkoxyimino group, or groups represented
by a formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p is any of a hydrogen
atom, or a lower alkyl group, a lower alkenyl group or a
lower alkynyl group which may be substituted with one to
10 three of said substituents, or a cyclo lower alkyl group,
an aryl group, or a heteroaromatic ring group selected from
a set of groups consisting of an imidazolyl group, an
isoxazolyl group, an isoquinolyl group, an isoindolyl group,
an indazolyl group, an indolyl group, an indoliziny group,
15 an isothiazolyl group, an ethylenedioxyphenyl group, an
oxazolyl group, a pyridyl group, a pyradinyl group, a
pyrimidinyl group, a pyridazinyl group, a pyrazolyl group,
a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl
group, a dihydroindolyl group, a thionaphthenyl group, a
20 naphthyridinyl group, a phenazinyl group, a benzoimidazolyl
group, a benzoxazolyl group, a benzothiazolyl group, a
benzotriazolyl group, a benzofuranyl group, a thiazolyl
group, a thiadiazolyl group, a thienyl group, a pyrrolyl
group, a furyl group, a furazanyl group, a triazolyl group,
25 a benzodioxanyl group and a methylenedioxyphenyl group, or
an aliphatic heterocyclic group(s) selected from a set of
groups consisting of an isoxazolinyl group, an
isoxazolidinyl group, a tetrahydropyridyl group, an
imidazolidinyl group, a tetrahydrofuranyl group, a

tetrahydropyranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, each of which(heteroaromatic
 5 ring groups and aliphatic heterocyclic groups) may be substituted with one to three of the same or different said substituent(s), which are same or different, or furthermore, may have (on it) a bicyclic or tricyclic fused ring of a partial structure selected from a set of groups consisting
 10 of:



; W_i is a single bond, an oxygen atom, a sulfur atom, SO , SO_2 , NR_q , SO_2NR_q , $N(R_q)SO_2NR_r$, $N(R_q)SO_2$, $CH(OR_q)$, $CONR_q$, $N(R_q)CO$, $N(R_q)CONR_r$, $N(R_q)COO$, $N(R_q)CSO$, $N(R_q)COS$, $C(R_q)=CR_r$,
 15 $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_q$, $OC(S)NR_q$, $SC(O)$, $SC(O)NR_q$ and $C(O)O$ (wherein: R_q and R_r are each, a hydrogen atom or a lower alkyl group, an aryl group or an aralkyl group, which may be substituted with one to three substituent(s) selected from a set of groups consisting of a lower alkyl
 20 group, a cyclo lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a
 25 carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower

- alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino
- 5 lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl
- 10 group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of said substituent(s).; Y_1 and Y_2 are each, the same or different, a single bond or a straight-chain or branched lower alkylene group which may have a
- 15 said bicyclic or tricyclic fused ring);
- 2) may have a five- to seven-membered fused ring selected from a set of groups consisting of:




- 20 which may be together with the carbon atom of said nitrogen-containing heteroaromatic cyclic group, on which the substituent, which is selected from a set of groups consisting of consisting of a lower alkyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower
- 25 alkyl group, a cyano lower alkyl group, a halo lower alkyl

group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, and a lower alkanoylamidino lower alkyl group (hereinafter indicated as ring-substituent) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent;

or,

3) may form a five- to seven-membered ring selected from a set of groups consisting of:



which may be formed from the carbon atom on which a substituent represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein: Y_1 , W_1 , Y_2 and R_p have the same meanings as stated above) stands, the carbon atom next to said carbon atom, and a

carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent; X and Z are each, the same or different, a carbon atom or a nitrogen atom, or being taken together with R₁ or R₂ and/or R₃ which may exist on X and Z, forms a

5 CH or a nitrogen atom; Y is CO, SO or SO₂;

R₁ is any of a hydrogen atom or a substituent represented by a formula Y₃-W₂-Y₄-R_s (wherein: R_s is any of a hydrogen atom or a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a cyclo lower alkyl group, an aryl group,

10 and a heteroaromatic ring group which is selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indoliziny group, an isothiazolyl group, an ethylenedioxyphenyl group, an

15 oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl

20 group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or

25 an aliphatic heterocyclic group selected from a set of groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl


11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, all of which may be substituted with one to three of said substituent(s); W_2 is a single bond, an oxygen atom, a
- 5 sulfur atom, SO , SO_2 , NR_t , SO_2NR_t , $N(R_t)SO_2NR_u$, $N(R_t)SO_2$, $CH(OR_t)$, $CONR_t$, $N(R_t)CO$, $N(R_t)CONR_u$, $N(R_t)COO$, $N(R_t)CSO$, $N(R_t)COS$, $C(R_v)=CR_x$, $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_t$, $OC(S)NR_t$, $SC(O)$, $SC(O)NR_t$ and $C(O)O$ (wherein: R_t and R_u are each a hydrogen atom or a substituent selected from a set of
 - 10 groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group,
 - 15 a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower
 - 20 alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower
 - 25 alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a lower alkyl group,



10031765-040203

- an aryl group or an aralkyl group which may be substituted with one to three of said substituent(s)); Y_3 and Y_4 are each, the same or different, a single bond or a straight-chain or branched lower alkylene group), or R_1 is a lower
- 5 alkyl group which may be substituted with one to three of the same or different substituent(s) which is selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a
- 10 lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower
- 15 alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower
- 20 alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl
- 25 group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a substituent or substituents selected from groups represented by the formula $Y_3-W_2-Y_4-R_5$ (wherein: R_5 , W_2 , Y_3 and Y_4 have the same meanings as stated

above), or R_1 forms a nitrogen atom together with X.); R_2 and R_3 are each independently, the same or different, a hydrogen atom, a hydroxy group, a lower alkyl group, a lower alkoxy group, or a substituent represented by the formula $Y_3-W_2-Y_4-R_5$ (wherein: R_5 , W_2 , Y_3 and Y_4 have the same meanings as stated above), or either one of R_2 or R_3 forms, together with R_1 and X, a saturated five- to eight-membered cyclic group selected from sets of groups of (a) and (b):



and the another one of R_2 or R_3 binds to a carbon atom or a nitrogen atom on the ring, or to a carbon atom, an oxygen atom and/or nitrogen atom on said ring-substituent to form a five- to seven-membered ring, or R_2 and R_3 are combined to form a spiro cyclo lower alkyl group, or are together furthermore with Z to which they bind to form an oxo (keto, or carbonyl) group, or they (R_2 and R_3) form, together with Z, R_1 and X, on which they stand, a saturated or an unsaturated five- to eight membered cyclic group which may be selected from sets of groups of (a) and (b):



- , which may contain one or more kinds of hetero atom(s)
- 5 selected from a group of a nitrogen atom, an oxygen atom and a sulfur atom, or which may be fused with any of a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group,
- 10 an isoindolyl group, an indazolyl group, an indolyl group, an indolydiny group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradiny group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxaliny group,
- 15 a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridiny group, a phenaziny group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl
- 20 group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazany group, a triazolyl group,

11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25

a benzodioxanyl group and a methylenedioxyphenyl group, or
 an aliphatic heterocyclic group(s) selected from a set of
 groups consisting of an isoxazolinyl group, an
 isoxazolidinyl group, a tetrahydropyridyl group, an
 5 imidazolidinyl group, a tetrahydrofuranyl group, a
 tetrahydropyranyl group, a piperazinyl group, a piperidinyl
 group, a pyrrolidinyl group, pyrrolinyl group, a morpholino
 group, a tetrahydroquinolinyl group and a
 tetrahydroisoquinolinyl group, which may be substituted
 10 with one to three of the same or different substituent(s)
 selected from a set of groups consisting of a lower alkyl
 group, a spiro cyclo lower alkyl group which may be
 substituted, a hydroxyl group, a cyano group, halogen atoms,
 a nitro group, a carboxyl group, a carbamoyl group, a
 15 formyl group, a lower alkanoyl group, a lower alkanoyloxy
 group, a hydroxy lower alkyl group, a cyano lower alkyl
 group, a halo lower alkyl group, a carboxy lower alkyl
 group, a carbamoyl lower alkyl group, lower alkoxy group, a
 lower alkoxycarbonyl group, lower alkoxycarbonylamino group,
 20 a lower alkoxycarbonylamino lower alkyl group, a lower
 alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a
 carbamoyloxy group, a lower alkylcarbamoyloxy group, di-
 lower alkylcarbamoyloxy group, an amino group, a lower
 alkylamino group, a di-lower alkylamino group, a tri-lower
 25 alkylammonio group, an amino lower alkyl group, a lower
 alkylamino lower alkyl group, a di-lower alkylamino lower
 alkyl group, a tri-lower alkylammonio lower alkyl group, a
 lower alkanoylamino group, an aroylamino group, a lower
 alkanoylamidino lower alkyl group, a lower alkylsulfinyl

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and a substituent or substituents selected from groups represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p , W_1 , Y_1 and Y_2 have the same meanings as stated above); R_4 and R_5 are each, the same or different, a hydrogen atom, halogen atoms, a hydroxy group, an amino group, or a substituent represented by the formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s , W_2 , Y_3 and Y_4 have the same meanings as stated above), or any of a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of the same or different substituent(s) selected from both a set of groups consisting of a lower alkyl group, a cyano group, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl

8. A method of preparing a compound of Formula (I) or pharmaceutically acceptable salts thereof:

$$\begin{array}{c}
 R_1 \quad R_2 \\
 \diagdown \quad \diagup \\
 X=Z \\
 \diagup \quad \diagdown \\
 Y \quad R_3 \\
 \diagdown \quad \diagup \\
 R_4 \quad R_5
 \end{array}
 \text{---} \text{HN} \begin{array}{c} \text{H} \\ \diagup \\ \text{C}=\text{O} \\ \diagdown \\ \text{N} \end{array} \text{---} \text{Ar} \quad (I)$$

15 (V) :


$$\begin{array}{c}
 R_{10} \quad R_{20} \\
 \diagdown \quad \diagup \\
 X = Z \\
 \diagup \quad \diagdown \\
 Y \quad R_{30} \\
 \diagdown \quad \diagup \\
 \text{---} \text{CON}_3 \text{ (V)} \\
 \diagup \quad \diagdown \\
 R_{40} \quad R_{50}
 \end{array}$$

20 with a compound represented with Formula (VI):

$$\text{H}_2\text{N}-\text{Ar}_0 \quad (\text{VI})$$

, wherein: Ar_0 have the same meanings as stated above,
to give a compound of Formula (II):


Formula (II)



5 , wherein: Ar_0 , X , Y , Z , R_{10} , R_{20} , R_{30} , R_{40} , R_{50} and the
formula --- have the same meanings as stated above
, and then, by removing, if necessary, the protective
group(s).


10 9. A method for preparing a compound of Formula (I')

Formula (I')




, wherein: T_1 is any of a single bond or a straight-chain
or branched lower alkylene, an aryl group, a heteroromatic
15 ring group, an aliphatic heterocyclic group, and an Ar
which has a convertible functional group(s) including or an
aralkyl group; Q is $W_1-Y_2-R_p$ (wherein: W_1 , Y_2 and R_p have the
same meanings as stated above), X , Y , Z , R_1 , R_2 , R_3 , R_4 , R_5
and the formula --- have the same meanings as stated above,
20 and salts thereof characterized by first making a compound
of a formula (VII):

Formula (VII)




- , wherein: L is a reactive substituent which may be protected, and may have a functional group which can be converted into other functional group, T₁₀ is any of a
- 5 single bond or , if appropriate, a straight-chain or branched lower alkylene group which may have a protected substituent(s), an aryl group, a heteroaromatic ring group, an aliphatic heterocyclic group, and an Ar₀ which has a convertible functional group including an aralkyl group,
- 10 reacting with a compound of a formula (VIII):

Formula (VIII)




- , wherein: R₆₀ is a hydrogen atom or a protective group for an amino group,
- 15 to obtain a compound of a formula (IX):

Formula (IX)




- , wherein: T₁₀, R₆₀ and L have the same meanings as stated above,
- 20 and then by making said compound react with a compound of a Formula (III):
- Formula (III)



- , wherein: X, Y, Z, R₁₀, R₂₀, R₃₀, R₄₀, R₅₀ and the formula
 --- have the same meanings as stated above] and one of
 reactive derivatives of formate ester, if necessary, in the
 5 presence of a base to give a compound of a formula (X):


Formula (X)



- , wherein: X, Y, Z, T₁₀, R₁₀, R₂₀, R₃₀, R₄₀, R₅₀, R₆₀ and the
 formula = have the same meanings as stated above,
 10 and by subjecting the compound obtained to transformation
 reaction of the substituent L and/or removal of the
 protective group.

10. A Cdk4 and/or Cdk6 inhibitory drug containing as the
 15 active ingredient a compound of Formula (I) and
 pharmaceutically acceptable salts thereof:

Formula (I)



, wherein: Ar is a nitrogen-containing heteroaromatic ring


100317561-0100000

- group selected from a set of groups consisting of a pyridyl group, a pyrimidinyl group, a pyradinyl group, a pyridazinyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyrazolyl group,
- 5 a pyrrolyl group, an imidazolyl group, an indolyl group, an isoindolyl group, a quinolyl group, an isoquinolyl group, a benzothiazolyl group, and a benzoxazolyl group, which:
- 1) may be substituted with one to three of the same or different substituent(s) selected from a set of groups
- 10 consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a
- 15 carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower
- 20 alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower
- 25 alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and groups represented

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

- by a formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p is any of a hydrogen atom, or a lower alkyl group, a lower alkenyl group or a lower alkynyl group which may be substituted with one to three of said substituents, or a cyclo lower alkyl group,
- 5 an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolizynyl group, an isothiazolyl group, an ethylenedioxyphenyl group, an
- 10 oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxalinyl group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl
- 15 group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or
- 20 an aliphatic heterocyclic group(s) selected from a set of groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a tetrahydropyranyl group, a piperazinyl group, a piperidinyl
- 25 group, a pyrrolidinyl group, pyrrolinyl group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, each of which cyclic group may be substituted with one to three of said substituents or, furthermore, may have (on it) a bicyclic or tricyclic


fused ring of a partial structure selected from a set of groups consisting of:



; W_1 is a single bond, an oxygen atom, a sulfur atom, SO,

- SO₂, NR_q, SO₂NR_q, N(R_q)SO₂NR_r, N(R_q)SO₂, CH(OR_q), CONR_q,
N(R_q)CO, N(R_q)CONR_r, N(R_q)COO, N(R_q)CSO, N(R_q)COS, C(R_q)=CR_r,
C≡C, CO, CS, OC(O), OC(O)NR_q, OC(S)NR_q, SC(O), SC(O)NR_q and
C(O)O (wherein: R_q and R_r are each a hydrogen atom or a
lower alkyl group, an aryl group or an aralkyl group, which
may be substituted with one to three substituent(s)
selected from a set of groups consisting of a lower alkyl
group, a cyclo lower alkyl group, a hydroxyl group, a cyano
group, halogen atoms, a nitro group, a carboxyl group, a
carbamoyl group, a formyl group, a lower alkanoyl group, a
lower alkanoyloxy group, a hydroxy lower alkyl group, a
cyano lower alkyl group, a halo lower alkyl group, a
carboxy lower alkyl group, a carbamoyl lower alkyl group,
lower alkoxy group, a lower alkoxycarbonyl group, lower
alkoxycarbonylamino group, a lower alkoxycarbonylamino
lower alkyl group, a lower alkylcarbamoyl group, a di-lower
alkylcarbamoyl group, a carbamoyloxy group, a lower
alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group,
an amino group, a lower alkylamino group, a di-lower
alkylamino group, a tri-lower alkylammonio group, an amino
lower alkyl group, a lower alkylamino lower alkyl group, a
di-lower alkylamino lower alkyl group, a tri-lower
alkylammonio lower alkyl group, a lower alkanoylamino group,
an aroylamino group, a lower alkanoylamidino lower alkyl

- group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and which may have one to three said substituent or substituents.); Y_1 and Y_2 are
- 5 each, the same or different, a single bond or a straight-chain or branched lower alkylene group which may have a said bicyclic or tricyclic fused ring);
- 2) may have a five- to seven-membered fused ring selected from a set of groups consisting of:




- which may be together with the carbon atom of said nitrogen-containing heteroaromatic ring group, on which the substituent, which is selected from a set of groups consisting of consisting of a lower alkyl group, a lower
- 15 alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower
- 20 alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-lower alkyl carbamoyloxy group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group,
- 25 an amino lower alkyl group, a lower alkylamino lower alkyl

group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, and
 5 a lower alkanoylamidino lower alkyl group (hereinafter indicated as ring-substituent) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent;

or,

- 10 3) may form a five- to seven-membered ring selected from a set of groups consisting of:



which may be formed from the carbon atom on which a substituent represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein:
 15 Y_1 , W_1 , Y_2 and R_p have the same meanings as stated above) stands, the carbon atom next to said carbon atom, and a carbon atom, an oxygen atom and/or a nitrogen atom on said ring-substituent.

; X and Z are each, the same or different, a carbon atom or
 20 a nitrogen atom, or being taken together with R_1 or R_2 and/or R_3 which may exist on X or Z, forms a CH or a nitrogen atom; Y is CO, SO or SO_2 ; R_1 is any of a hydrogen atom or a substituent represented by a formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s is a hydrogen atom or a lower alkyl group, a
 25 lower alkenyl group, a lower alkynyl group, a cyclo lower


alkyl group, an aryl group, and a heteroaromatic ring group selected from a set of groups consisting of an imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indoliziny group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradinyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazolyl group, a quinoxaliny group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridinyl group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group (which is) selected from a set of groups consisting of an isoxazoliny group, an isoxazolidiny group, a tetrahydropyridyl group, an imidazolidiny group, a tetrahydrofuranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrroliny group, a morpholino group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, all of which may be substituted with one to three of said substituent(s); W_2 is a single bond, an oxygen atom, a sulfur atom, SO , SO_2 , NR_t , SO_2NR_t , $N(R_t)SO_2NR_u$, $N(R_t)SO_2$, $CH(OR_t)$, $CONR_t$, $N(R_t)CO$, $N(R_t)CONR_u$, $N(R_t)COO$, $N(R_t)CSO$, $N(R_t)COS$, $C(R_v)=CR_x$, $C\equiv C$, CO , CS , $OC(O)$, $OC(O)NR_t$, $OC(S)NR_t$, $SC(O)$, $SC(O)NR_t$ and $C(O)O$ (wherein: R_t and R_u are each

4003756-030000


- hydrogen atom or a substituent selected from a set of groups consisting of a lower alkyl group, a hydroxy group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of said substituent or substituents); Y_3 and Y_4 are each, the same or different, a single bond or a straight-chain or branched lower alkylene group, or R_1 is an lower alkyl group which may be substituted with one to three of the same or different substituent or substituents selected from a set of groups consisting of a lower alkyl group, a hydroxyl group, a cyano group, halogen atoms, a

40031781-01-000000


- nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a
- 5 carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxy carbonyl group, lower alkoxy carbonylamino group, a lower alkoxy carbonylamino lower alkyl group, a lower alkyl carbamoyl group, a di-lower alkyl carbamoyl group, a carbamoyloxy group, a lower alkyl carbamoyloxy group, di-
- 10 lower alkyl carbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkyl ammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkyl ammonio lower alkyl group, a
- 15 lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and a substituent selected from groups
- 20 represented by the formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s , W_2 , Y_3 and Y_4 have the same meanings as stated above), or R_1 forms a nitrogen atom together with X; R_2 and R_3 are each independently, the same or different, a hydrogen atom, a hydroxy group, a lower alkyl group, a lower alkoxy group,
- 25 or a substituent represented by the formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s , W_2 , Y_3 and Y_4 have the same meanings as stated above), or either one of R_2 or R_3 forms, together with R_1 and X, a saturated five- to eight-membered cyclic group selected from sets of groups of (a) and (b):




and



- and the other one of R_2 or R_3 binds to a carbon atom or a
 5 nitrogen atom on the ring, or to a carbon atom, an oxygen
 atom and/or nitrogen atom on said ring-substituent to form
 a five- to seven-membered ring, or R_2 and R_3 are combined
 to form a spiro cyclo lower alkyl group, or R_2 and R_3 are
 combined furthermore with Z on which they exist to form an
 10 oxo (keto, or carbonyl) group, or they (R_2 and R_3) form,
 together with Z, R_1 and X on which they stand, a saturated
 or an unsaturated five- to eight-membered cyclic group
 which may be selected from sets of groups of (a) and (b):



15 and



which may contain one or more kinds of hetero atom(s)

selected from a group of a nitrogen atom, an oxygen atom and a sulfur atom, or which may be fused with any of a cyclo lower alkyl group, an aryl group, a heteroaromatic ring group selected from a set of groups consisting of an

5 imidazolyl group, an isoxazolyl group, an isoquinolyl group, an isoindolyl group, an indazolyl group, an indolyl group, an indolydiny group, an isothiazolyl group, an ethylenedioxyphenyl group, an oxazolyl group, a pyridyl group, a pyradiny group, a pyrimidiny group, a

10 pyridazinyl group, a pyrazolyl group, a quinoxaliny group, a quinolyl group, a dihydroisoindolyl group, a dihydroindolyl group, a thionaphthenyl group, a naphthyridiny group, a phenazinyl group, a benzoimidazolyl group, a benzoxazolyl group, a benzothiazolyl group, a

15 benzotriazolyl group, a benzofuranyl group, a thiazolyl group, a thiadiazolyl group, a thienyl group, a pyrrolyl group, a furyl group, a furazanyl group, a triazolyl group, a benzodioxanyl group and a methylenedioxyphenyl group, or an aliphatic heterocyclic group(s) selected from a set of

20 groups consisting of an isoxazolinyl group, an isoxazolidinyl group, a tetrahydropyridyl group, an imidazolidinyl group, a tetrahydrofuranyl group, a tetrahydropyranyl group, a piperazinyl group, a piperidinyl group, a pyrrolidinyl group, pyrrolinyl group, a morpholino

25 group, a tetrahydroquinolinyl group and a tetrahydroisoquinolinyl group, which may be substituted with one to three of the same or different substituent(s) selected from a set of groups consisting of a lower alkyl group, a spiro cyclo lower alkyl group which may be

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

substituted, a hydroxyl group, a cyano group, halogen atoms, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, and a substituent selected from groups represented by the formula $Y_1-W_1-Y_2-R_p$ (wherein: R_p , W_1 , Y_1 and Y_2 have the same meanings as stated above); R_4 and R_5 are each, the same or different, a hydrogen atom, halogen atoms, a hydroxy group, an amino group, or a substituent represented by the formula $Y_3-W_2-Y_4-R_s$ (wherein: R_s , W_2 , Y_3 and Y_4 have the same meanings as stated above), or any of a lower alkyl group, an aryl group or an aralkyl group which may be substituted with one to three of the same of different substituent or substituents selected from both a

10034735-040200

set of groups consisting of a lower alkyl group, a cyano group, a nitro group, a carboxyl group, a carbamoyl group, a formyl group, a lower alkanoyl group, a lower alkanoyloxy group, a hydroxy lower alkyl group, a cyano lower alkyl group, a halo lower alkyl group, a carboxy lower alkyl group, a carbamoyl lower alkyl group, lower alkoxy group, a lower alkoxycarbonyl group, lower alkoxycarbonylamino group, a lower alkoxycarbonylamino lower alkyl group, a lower alkylcarbamoyl group, a di-lower alkylcarbamoyl group, a carbamoyloxy group, a lower alkylcarbamoyloxy group, di-lower alkylcarbamoyloxy group, an amino group, a lower alkylamino group, a di-lower alkylamino group, a tri-lower alkylammonio group, an amino lower alkyl group, a lower alkylamino lower alkyl group, a di-lower alkylamino lower alkyl group, a tri-lower alkylammonio lower alkyl group, a lower alkanoylamino group, an aroylamino group, a lower alkanoylamidino lower alkyl group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfonylamino group, a hydroxyimino group and a lower alkoxyimino group, or groups represented by the formula $Y_3-W_2-Y_4-R_8$ (wherein: R_8 , W_2 , Y_3 and Y_4 have the same meanings as stated above); and the formula \equiv represents either a single bond or a double bond.